

Family Planning, Access, Care, and Treatment Clinical Services Benefit Grid

The following information, referenced from the Family PACT Clinical Services Benefit Grid (Revised February 2020), serves as a reference tool for laboratory test services and is not comprehensive. For comprehensive lists of diagnosis codes and procedures, please refer to the appropriate sections in the Family PACT Clinical Services and Billing Instructions Manual. As a reminder, medical record documentation is required to support all lab tests that are requested. The Family PACT program benefits state: Medical record documentation must include of providing, ordering, or deferring services rendered to clients, according to the *Program Standards* section, including, but not limited to, client assessment, diagnosis, treatment, and follow-up. Medical record documentation must include

					Fema	le				Ma	ile and Fem	ale	<u> </u>	Male	_	
Blank = Test(s)	vered under ICD-10-CM code. not covered. additional diagnosis code. See reverse side of grid for family planning-	Emergency contraceptive counseling. Procreative counseling using NFP. General contraception management counseling and advice.	Initiation of contraceptive pills. Initiation of vaginal ring. Initiation of transdermal patch.	Surveillance of contraceptive pills. Surveillance of vaginal ring usage. Surveillance of transdermal patch.	initiation of contraceptive injection. Insertion of implantable subdermal contraceptive.	Survelliance of contraceptive injection. Surveillance of implantable subdermal contraceptive (checking, reinsertion, or removal)	Insertion of intrauterine contraceptive device (IUC). Surveiliance of IUC.	Removal of IUC. Removal and reinsertion of IUC.	Sterilization procedure. Tubal ligation status.	Initiation of other contraceptive methods: barriers, diaphragm. Natural family planning method.	Surveillance of other contraceptive methods: barriers, diaphragm.	Pre-procedural lab exam. Sterilization counseling and advice.	Sterilization procedure. Vasectomy status.	Postprocedural hemorrhage of a genitourinary system organ or structure following a genitourinary system procedure		For specific indications of diagnostic testing for HPV, CT, and NG and for complication services and corresponding diagnosis codes, please refer to the Family PACT Policies, Procedures, and Billing Instructions Manual, available at www.medical.ca.gov.
Quest Diagnostics Order Code	Family Planning ICD-10-CM Diagnosis Codes EVERY LABORATORY REQUEST MUST INCLUDE A CONTRACEPTIVE METHOD.	Z30.012 Z31.61 Z30.09	Z30.011 Z30.015 Z30.016	Z30.41 Z30.44 Z30.45	Z30.013 Z30.017	Z30.42 Z30.46	Z30.430 Z30.431	Z30.432 Z30.433	Z30.2 Z98.51	Z30.018 Z30.02	Z30.49	Z01.812 AND Z30.09	Z30.02 Z98.52 Z30.018 Z30.49	N99.820		
	Family PACT Laboratory Services														Fami	ily Planning-Related Services
1759	Auto CBC w/out differential (1) (2)								√ (2)			√ (1)				
6399	CBC w/ Platelets (1) (2) (3)		SEC	SEC	SEC	SEC	SEC	SEC	√ (2)	SEC (F)	✓SEC (F)	√ (1)	√ (3)			tory tests listed below are FPACT-Related Services ra tests may require an additional diagnosi code.
11361	Chlamydia trachomatis RNA, TMA (4)		√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)		√ (4)			of Grid for approved ICD 10-CM diagnosis codes.
11363	CT/NG RNA, TMA (4)		√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)		√ (4)		Order Code	Family Planning-Related
11362	Neisseria gonorrhoeae RNA, TMA (4)		√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)	√ (4)		√ (4)		11361	Laboratory Services Chlamydia Trachomatis, RNA
396	HCG Qualitative, Urine	✓	7	✓	√	1	✓	1	7	✓	√				11362	Neisseria Gonorrhoeae, RNA
509	Hematocrit						-	√	√ (2)			√ (1)		1	6399	CBC w/Diff
510	Hemoglobin						~	1	√ ₍₂₎			√ (1)			809 497	ESR Sed Rate Auto Gram Stain (male only)
91431	HIV-1/HIV-2 Antigen/Antibody with reflexes		7	7	7	√	7	1	7	✓	✓		√			or 90887 HPV HR (female only)
Reflex	HPV Genotypes 16, 18, and 45 (if performed) (7)		✓SEC	✓SEC	√SEC	√SEC	✓SEC	√SEC	✓SEC	✓SEC (F)	✓SEC (F)				Reflex	HPV Genotype 16, 19/45 (female only)
31532 (DNA) or 90887 (mRNA)	HPV Screening, reflexive/concurrent (5)		✓SEC	✓SEC	✓SEC	✓SEC	✓SEC	√SEC	✓SEC	✓SEC	✓SEC				2692 34257,	HSV Culture 90569, 90570 HSV 1 & 2 NAAT
	8810, 18811, 58315, 90934, 92087, 92236, 92238, 92245, 92094, 91414 PAP Smear, Thin Prep®, or Sure Path™ (6)		1	1	1	1	1	1	1	1	1				5367 799	PH (female only) Syphylis - Quantitive
3542	Surgical Pathology								✓				1		3542	Surgical Pathology
4112 (IFA) or	Treponemal pallidum (TP) confirmation (7)				1					1	_				19550	or 90521 Trichomonas vaginalis, RNA, TMA (females only)
653 (PA)	immunofluorescent assay or particle agglutination		✓	✓	•	✓	✓	✓	*		V		✓		8563	Urine Microscopy*
5463	Urinalysis with Microscopic (1)		SEC	SEC	SEC	SEC	SEC	SEC	SEC	SEC (F)	SEC (F)	√ (1)			5463	Urinalysis w/ Microscopy*
6448	Urinalysis without Microscopic (1)		SEC	SEC	SEC	SEC	SEC	SEC	SEC	SEC (F)	SEC (F)	√ ₍₁₎			6448	Urinalysis w/o Microscopy* 4 Wet Mount
799 or 36126	VDRL, RPR		-	7	√	√	-	✓	~	~	~		✓		91475	Mycoplasma genitalium
(1) Pre-operative	testing only. For sterilization services both Z01.812 and Z30.0	9 are required														* Symptom(s) suggestive of UTI

- (2) Postoperative testing only with female sterilization code Z98.51.
- (3) Evaluation of postoperative infection with male sterilization code Z98.52.
- (4) Females less than 25 years of age: routine annual screening covered under primary family planning ICD-10-CM code.

Age and gender restrictions apply to more frequent screening and to diagnostic testing. See reverse for additional screening and diagnostic ICD-10-CM codes.

- (5) Limited to reflex testing from ASC-US cytology for women ≥ 21; concurrent testing (co-testing) based on medical necessity and age. Co-testing for primary screening covered every 5 years for women ≥ 30 with additional ICD-10 code Z11.51. See reverse for additional ICD-10-CM codes required for co-testing.
- (6) Routine screening restricted to women ages 21 through 65.
- (7) Limited to reflex testing subsequent to a positive test result, and only reimbursable with or after a paid claim.

Diagnosis must always be documented in the patient's medical record. The ultimate responsibility belongs to the ordering physician to correctly assign the patient's diagnosis based on the patient's history, symptoms, and medical condition.

Note: Complication services require a TAR (Treatment Authorization Request form). See Family PACT Policies, Procedures, and Billing Instructions Manual or website for more information regarding complication services and corresponding ICD-10-CM codes.

Family Planning-Related Services

Family Planning ICD-10-CM Diagnosis Code Required for Every Claim

An additional ICD-10-CM diagnosis code is required as indicated on tests listed below.

6399	CBC/diff (not required for Z30.09, Z30.2					
0399	Z98.51, Z98.52					
809	ESR (sed rate) automated					
29891	ESR (sed rate) non-automated					
N70.03 (F)	Acute salpingitis and oophoritis					
N70.93 (F)	Salpingitis and oophoritis, unspecified					
N94.10 (F)	Unspecified dyspareunia					
N94.11 (F)	Superficial (introital) dyspareunia					
N94.12 (F)	Deep dyspareunia					
N94.19 (F)	Other specified dyspareunia					
N94.89 (F)	Other conditions associated with female genital organs and menstrual cycle					

799	Syphilis - Quantitative
A51.0 (M/F)	Primary genital syphilis
A51.31 (M/F)	Condyloma latum
A51.39 (M/F)	Other secondary syphilis of skin
A51.5 (M/F)	Early syphilis, latent
A52.8 (M/F)	Late syphilis, latent
A53.0 (M/F)	Latent syphilis, unspecified

2692	HSV Culture
34257, 90569, 90570	HSV 1 & 2 NAAT
N48.5 (M)	Ulcer of penis
N76.6 (F)	Ulceration of vulva
HSV typing is not	covered

11361 (CT)	Chlamydia Trachomatis and Neisseria					
11362 (NG)	Gonorrhoeae					
or 11363						
Z11.63 (M/F)	Screening for bacterial STIs (NG)					
Z11.8 (M/F)	Screening for other infections (CT)					
Z20.2 (M/F)	Contact with or exposure to sexually transmitted infections					
Z22.4 (M/F)	Carrier or suspected carrier of STIs					
Z72.51 (M/F)	High-risk heterosexual behavior					
Z72.52 (M/F)	High-risk homosexual behavior					
Z72.53 (M/F)	High-risk bisexual behavior					
Z86.19 (M/F)	Personal history of other diseases					

91475	Mycoplasma Genitalium
N70.03 (F)	Acute salpingitis and oophoritis
N70.93 (F)	Salpingitis and oophoritis, unspecified
N72 (F)	Inflammatory disease of cervix uteri
N34.1 (M)	Nonspecific urethritis
N34.2 (M)	Other urethritis
N34.3 (M)	Urethral syndrome, unspecified

5463	Urinalysis, dipstick or auto w/ micro
6448	Urinalysis, dipstick or auto w/o micro, Urine
8563	Micrscopy
N30.00 (F)	Acute cystitis without hematuria
N30.01 (F)	Acute cystitis with hematuria
R10.30 (F)	Lower abdominal pain, unspecified
R30.0 (F)	Dysuria
R30.9 (F)	Painful micturition, unspecified
R31.0 (F)	Gross hematuria
R35.0 (F)	Frequency of micturition
N34.1 (M)	Nonspecific urethritis
N45.1 (M)	Epididymitis
N45.3 (M)	Epididymo-orchitis
N50.811 (M)	Right testicular pain
N50.812 (M)	Left testicular pain
N50.819 (M)	Testicular pain, unspecified

497	Gram Stain (males only) CT and NG symptomatic
A54.01 (M)	Gonococcal cystitis and urethritis
A54.22 (M)	Gonococcal prostatitis
A54.5 (M)	Gonococcal pharyngitis
A54.6 (M)	Gonococcal infection, anus/rectum
A56.01 (M)	Chlamydial cystitis and urethritis
A56.3 (M)	Chlamydial infection, anus/rectum
N34.1 (M)	Nonspecific urethritis
N34.2 (M)	Other urethritis
N45.3 (M)	Epididymo orchitis
N45.1 (M)	Epididymitis
N50.811 (M)	Right testicular pain
N50.812 (M)	Left testicular pain
N50.819 (M)	Testicular pain, unspecified

11361 (CT), 113	362 (NG), Chlamydia	Trachomatis and Ne	eisseria Gonorrhoeae
or 11363		Additional ICD-10-C	M Codes
	Males and Females		Females Only
A54.01(M/F)	Gonococcal cystitis and urethritis, unspecified	A54.03 (F)	Gonococcal cervicitis
A54.5 (M/F)	Gonococcal pharyngitis	N70.03 (F)	Acute salpingitis and oophoritis
A54.6 (M/F)	Gonococcal infection of anus and rectum	N70.93 (F)	Salpingitis and oophoritis, unspecified
A56.01 (M/F)	Chlamydial cystitis and urethritis	N72 (F)	Inflammatory disease of cervix uteri
A56.3 (M/F)	Chlamydial infection of anus and rectum	N89.8 (F)	Vaginal leukorrhea
A56.4 (M/F)	Chlamydial infection of pharynx	N94.89 (F)	Other conditions assoc with female
R30.0 (M/F)	Dysuria	N94.10-N94.19 (F)	Dyspareunia
R30.9 (M/F)	Painful micturition, unspecified		Males Only
Z20.2 (M/F)	Contact with or exposure to sexually transmitted	A54.22 (M)	Gonococcal prostatitis
220.2 (W/F)	infections	N34.1 (M)	Nonspecific urethritis
		N34.2 (M)	Other urithritis
		N45.3 (M)	Epididymo-orchitis

31532 90887 Females only (age guide)	HPV-amplified probe high-risk (limited to one per year per client) females age 21-99. Additional ICD-10-CM code required beyond ASC-US reflex testing based on age.
D06.9 (≥21)	Carcinoma in situ of cervix
N87.0 (≥25)	Mild cervical dysplasia
N87.1 (≥21)	Moderate cervical dysplasia
R87.610 (≥21)	ASC-US Pap
R87.611 (≥25) R87.612 (≥25)	ASC-H Pap LGSIL Pap
R87.613 (≥25)	HGSIL Pap
R87.616 (≥30)	Satisfactory cervical smear but lacking transformation zone
R87.619 (≥21)	Unspecified abnormal cytological findings, cervix
Z01.42 (≥21)	Pap to confirm normal after abnormal smear
Z87.410 (≥21)	Personal history of cervical dysplasia

509	Hematocrit
Z30.2 (M)	Encounter for sterilization
Z30.09 (F)	Encounter for other general counseling and advice on contraception
Z30.42 (F)	Encounter for surveillance of injectable contraceptive

LAC92214	Wet mount (gel swab)
A59.01 (F)	Trichomonal vulvovaginitis
B37.3 (F)	Candidal vulvovaginitis
N34.1 (M)	Nonspecific urethritis
A59.03 (M/F)	Trichomonal cystitis and urethritis
N34.2 (M)	Other urithritis
N76.0 (F)	Acute vaginitis
Z20.2 (M/F)	Trichomoniasis-exposed partner

3542	Surgical Pathology
A63.0 (M/F)	Anogenital (venereal) warts
B07.9 (M/F)	Viral wart, unspecified
B08.1 (M/F)	Molluscum contagiosum
D06.9 (F)	Carcinoma in situ of cervix
N87.0 (F)	Mild cervical dysplasia
N87.1 (F)	Moderate cervical dysplasia
N88.0 (F)	Leukoplakia of cervix uteri
R87.610 (F)	ASC-US cervical cytology
R87.611 (F)	ASC-H cervical cytology
R87.612 (F)	LGSIL cervical cytology
R87.613 (F)	HGSIL cervical cytology
R87.616 (F)	Satisfactory cervical smear but lacking transformation zone
R87.618 (F)	Other abnormal cytological findings, cervix
R87.619 (F)	Unspecified abnormal cytological findings, cervix
R87.810 (F)	Cervical high-risk HPV positive

3542	Surgical Pathology, LEEP only; females age ≥15 years
D06.9 (F)	Carcinoma in situ of cervix
N87.0 (F)	Mild cervical dysplasia
N87.1 (F)	Moderate cervical dysplasia

Reflex	HPV Genotypes 16, 18/45 females age 30-65 years
Reflexive from	a positive hrHPV test
R87.810 (F)	Cervical high risk human papillomavirus (HPV) test positive

19550 or 90521	Thricomonas vaginalis, RNA, TMA Qualitative (female only)
A59.01 (F)	Trichomonal vulvovaginitis
A59.03 (F)	Trichomonal cystitis and urethritis
N76.0 (F)	Acute vaginitis
Z20.2 (F)	Trichomoniasis-exposed partner

For specific indications of diagnostic testing for HPV, CT and NG, and for complication services and corresponding diagnosis codes, please refer to the Family Pact Policies, Procedures, and Billing Instructions Manual, available at medi-cal.ca.gov

Updated March 2023