




Patient Information	Specimen Information	Client Information
TEST, PATIENT DOB: 01/01/1965 AGE: 50 Gender: F Fasting: Y Phone: NG Patient ID: BH047903A	Specimen: BH047903A Requisition: 0422196 Lab Ref #: 0422196 Collected: 04/07/2015 / 11:28 EDT Received: 04/07/2015 / 11:29 EDT Reported: 04/07/2015 / 14:13 EDT	Client #: 97502840 AARRSSS COLMENAR, ANTONIO TEST CLIENT (HQ) AH1 1 MALCOLM AVE TETERBORO, NJ 07608-1011

COMMENTS: TEST PATIENT


Liver Fibrosis, FibroTest™ ActiTest™ Panel

Fibro Test
Fibro Test assesses the fibrosis of the liver



Score : 0.62
(F3)
advanced fibrosis

Acti Test
Acti Test assesses activity (inflammation in chronic viral hepatitis C or B)



Score : 0.18
(A0-A1)
no activity

Biomarkers

Lab: EZ

Analyte	Result	Reference Range/Comments	Analyte	Result	Reference Range/Comments
Alpha2 Macroglobulin	301 H	106-279 mg/dL	Total Bilirubin	0.4	0.2-1.2 mg/dL
Haptoglobin	47	43-212 mg/dL	Gamma GT	63	3-70 U/L
Apolipoprotein A1	104	101-198 mg/dL	ALT	26	6-29 U/L

Liver Fibrosis History

Previous Results

No historical results currently available.

Only includes results that meet strict identification matching.
Historical values represent Fibrotests performed at Quest Diagnostics only; previous results from tests performed at other laboratories have not been included.

Reference ID for internal report tracking 14072

Lab: EZ

Precautions of use and interpretability

Lab: EZ

- The reliability of results is dependent on compliance with the preanalytical and analytical conditions recommended by BioPredictive.
- The Tests have to be deferred for: acute hemolysis, acute hepatitis, acute inflammation, extra hepatic cholestasis.
- The advice of a specialist should be sought for interpretation in chronic hemolysis and Gilbert's syndrome.
- The Test interpretation is not validated in liver transplant patients.
- Isolated extreme values of one of the components should lead to caution in interpreting the results.
- In case of discordance between a biopsy result and a Test, it is recommended to seek the advice of a specialist. The causes of these discordances could be due to a flaw of the Test or to a flaw in the biopsy: i.e. a liver biopsy has a 33% variability rate for one fibrosis stage
- FibroTest is interpretable for chronic hepatitis B and C, alcoholic and non alcoholic steatosis.
- ActiTest is interpretable for chronic hepatitis B and C.

The performance characteristics have been determined by Quest Diagnostics Nichols Institute, San Juan Capistrano. It has not been cleared or approved by the U.S. Food and Drug Administration. Performance characteristics refer to the analytical performance of the test.

