

## Family Planning, Access, Care, and Treatment Clinical Services Benefit Grid

The following information, referenced from the Family PACT Clinical Services Benefit Grid (Revised March 2024), serves as a reference tool for laboratory test services and is not comprehensive lists of diagnosis codes and procedures, please refer to the appropriate sections in the Family PACT Policies, Procedures, and Billing Instructions Manual. As a reminder, medical record documentation is required to support claims for providing, ordering, or deferring services rendered to clients, according to the *Program Standards* section, including, but not limited to, client assessment, diagnosis, treatment, and follow-up. Medical record documentation must include justification to support claims for providing and providing and

					Fema	le				Ma	ale and Fem	ale	Male
Grid Key  ☑ = Test(s) covered under ICD-10-CM code.  Blank = Test(s) not covered.  SEC = Requires additional diagnosis code. See reverse side of grid for family planning-related services.		Emergency contraceptive counseling. Procreative counseling using NFP. General contraception management counseling and advice.	initiation of contraceptive pills. Initiation of vaginal ring. Initiation of transdermal patch.	Surveillance of contraceptive pills. Surveillance of vaginal ring usage. Surveillance of transdermal patch.	initiation of contraceptive injection. Insertion of implantable subdermal contraceptive.	Survelliance of contraceptive injection. Surveillance of implantable subdermal contraceptive (checking, reinsertion, or removal)	Insertion of intrauterine contraceptive device (IUC). Surveiliance of IUC.	Removal of IUC. Removal and reinsertion of IUC.	Sterilization procedure. Tubal ligation status.	initiation of other contraceptive methods: barriers, diaphragm. Natural family planning method.	Surveillance of other contraceptive methods: barriers, diaphragm.	Pre-procedural lab exam. Sterilization counseling and advice.	Sterilization procedure. Vasectomy status.
Quest Diagnostics Order Code	Family Planning ICD-10-CM Diagnosis Codes EVERY LABORATORY REQUEST MUST INCLUDE A CONTRACEPTIVE METHOD.	Z30.012 Z31.61 Z30.09	Z30.011 Z30.015 Z30.016	Z30.41 Z30.44 Z30.45	Z30.013 Z30.017	Z30.42 Z30.46	Z30.430 Z30.431	Z30.432 Z30.433	Z30.2 Z98.51	Z30.018 Z30.02	Z30.49	Z01.812 AND Z30.09	Z30.02 Z98.52 Z30.018 Z30.49
	Family PACT Laboratory Services												
1759	Auto CBC w/out differential (1) (2)								<b>√</b> (2)			<b>√</b> (1)	
6399	CBC w/ Platelets (1) (2) (3)		SEC	SEC	SEC	SEC	SEC	SEC	<b>√</b> (2)	SEC (F)	✓SEC (F)	<b>√</b> (1)	<b>√</b> (3)
11361	Chlamydia trachomatis RNA, TMA (4)		<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)		<b>√</b> (4)
11363	CT/NG RNA, TMA (4)		<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)		<b>√</b> (4)
11362	Neisseria gonorrhoeae RNA, TMA (4)		<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)	<b>√</b> (4)		<b>√</b> (4)
396	HCG Qualitative, Urine	✓	<b>-</b>	<b>1</b>	<b>✓</b>	<b>*</b>	<b>1</b>	✓	<b>-</b>	<b>-</b>	<b>V</b>		
509	Hematocrit						<b>-</b>	✓	<b>√</b> (2)			<b>√</b> (1)	
510	Hemoglobin						<b>*</b>	✓	<b>√</b> (2)			<b>√</b> (1)	
91431	HIV-1/HIV-2 Antigen/Antibody with reflexes		<b>—</b>	<b>-</b>	<b>✓</b>	✓	<b>-</b>	✓	<b>-</b>	<b>—</b>			✓
Reflex	HPV Genotypes 16, 18, and 45 (if performed) (7)		✓SEC	✓SEC	✓SEC	✓SEC	✓SEC	✓SEC	✓SEC	✓SEC (F)	✓SEC (F)		
31532 (DNA) or 90887 (mRNA)	HPV Screening, reflexive/concurrent (5)		✓SEC	✓SEC	✓SEC	✓SEC	✓SEC	√SEC	✓SEC	✓SEC	✓SEC		
	18810, 18811, 58315, 90934, 92087, 92236, 92238, 92245, 8, 92094, 91414 PAP Smear, Thin Prep®, or Sure Path™ (6)		1	<b>√</b>	✓	1	<b>√</b>	1	1	1	<b>1</b>		
3542	Surgical Pathology								7				✓
4112 (IFA) or 653 (PA)	Treponemal pallidum (TP) confirmation (7) immunofluorescent assay or particle agglutination		1	1	1	<b>√</b>	1	1	1	1	1		<b>√</b>
5463	Urinalysis with Microscopic (1)		SEC	SEC	SEC	SEC	SEC	SEC	SEC	SEC (F)	SEC (F)	<b>√</b> (1)	
6448	Urinalysis without Microscopic (1)		SEC	SEC	SEC	SEC	SEC	SEC	SEC	SEC (F)	SEC (F)	<b>√</b> (1)	

For specific indications of diagnostic testing for HPV, CT, and NG and for complication services and corresponding diagnosis codes, please refer to the Family PACT Policies, Procedures, and Billing Instructions Manual, available at www.medical.ca.gov.

Family Planning-Related Services

The laboratory tests listed below are FPACT-Related Services. These labora tests may require an additional diagnosi code. See back of Grid for approved ICD 10-CM diagnosis codes.

Order	Family Planning-Related
Code	Laboratory Services
11361	Chlamydia Trachomatis, RNA
11362	Neisseria Gonorrhoeae, RNA
6399	CBC w/Diff
809	ESR Sed Rate Auto
497	Gram Stain (male only)
31532 or	90887 HPV HR (female only)
D (1	HPV Genotype 16, 19/45
Reflex	(female only)
2692	HSV Culture
34257, 9	0569, 90570 HSV 1 & 2 NAAT
5367	PH (female only)
799	Syphylis - Quantitive
3542	Surgical Pathology
19550 or	90521 Trichomonas vaginalis,
	RNA, TMA (females only)
8563	Urine Microscopy*
5463	Urinalysis w/ Microscopy*
6448	Urinalysis w/o Microscopy*
LAC92214	Wet Mount
91475	Mycoplasma genitalium
	* Symptom(s) suggestive of UTI

- (1) Pre-operative testing only. For sterilization services both Z01.812 and Z30.09 are required.
- (2) Postoperative testing only with female sterilization code Z98.51.
- (3) Evaluation of postoperative infection with male sterilization code Z98.52.
- (4) Females less than 25 years of age: routine annual screening covered under primary family planning ICD-10-CM code.

Age and gender restrictions apply to more frequent screening and to diagnostic testing. See reverse for additional screening and diagnostic ICD-10-CM codes.

- (5) Limited to reflex testing from ASC-US cytology for women ≥ 21; concurrent testing (co-testing) based on medical necessity and age. Co-testing for primary screening covered every 5 years for women ≥ 30 with additional ICD-10 code Z11.51. See reverse for additional ICD-10-CM codes required for co-testing.
- (6) Routine screening restricted to women ages 21 through 65.
- (7) Limited to reflex testing subsequent to a positive test result, and only reimbursable with or after a paid claim.

Diagnosis must always be documented in the patient's medical record. The ultimate responsibility belongs to the ordering physician to correctly assign the patient's diagnosis based on the patient's history, symptoms, and medical condition.

Note: Complication services require a TAR (Treatment Authorization Request form). See Family PACT Policies, Procedures, and Billing Instructions Manual or website for more information regarding complication services and corresponding ICD-10-CM codes.

## **Family Planning-Related Services**

## Family Planning ICD-10-CM Diagnosis Code Required for Every Claim

An additional ICD-10-CM diagnosis code is required as indicated on tests listed below.

6399	CBC/diff (not required for Z30.09, Z30.2
6399	Z98.51, Z98.52
809	ESR (sed rate) automated
29891	ESR (sed rate) non-automated
N70.03 (F)	Acute salpingitis and oophoritis
N70.93 (F)	Salpingitis and oophoritis, unspecified
N94.10 (F)	Unspecified dyspareunia
N94.11 (F)	Superficial (introital) dyspareunia
N94.12 (F)	Deep dyspareunia
N94.19 (F)	Other specified dyspareunia
N94.89 (F)	Other conditions associated with female genital organs and menstrual cycle

799	Syphilis - Quantitative
A51.0 (M/F)	Primary genital syphilis
A51.31 (M/F)	Condyloma latum
A51.39 (M/F)	Other secondary syphilis of skin
A51.5 (M/F)	Early syphilis, latent
A52.8 (M/F)	Late syphilis, latent
A53.0 (M/F)	Latent syphilis, unspecified

2692	HSV Culture		
34257, 90569, 90570	HSV 1 & 2 NAAT		
N48.5 (M)	Ulcer of penis		
N76.6 (F)	Ulceration of vulva		
HSV typing is not covered			

11361 (CT)	Chlamydia Trachomatis and Neisseria				
11362 (NG)	Gonorrhoeae				
or 11363					
Z11.63 (M/F)	Screening for bacterial STIs (NG)				
Z11.8 (M/F)	Screening for other infections (CT)				
Z20.2 (M/F)	Contact with or exposure to sexually				
220.2 (141/1 )	transmitted infections				
Z22.4 (M/F)	Carrier or suspected carrier of STIs				
Z72.51 (M/F)	High-risk heterosexual behavior				
Z72.52 (M/F)	High-risk homosexual behavior				
Z72.53 (M/F)	High-risk bisexual behavior				
Z86.19 (M/F)	Personal history of other diseases				

91475	Mycoplasma Genitalium
N70.03 (F)	Acute salpingitis and oophoritis
N70.93 (F)	Salpingitis and oophoritis, unspecified
N72 (F)	Inflammatory disease of cervix uteri
N34.1 (M)	Nonspecific urethritis
N34.2 (M)	Other urethritis
N34.3 (M)	Urethral syndrome, unspecified

5463	Urinalysis, dipstick or auto w/ micro
6448	Urinalysis, dipstick or auto w/o micro, Urine
8563	Micrscopy
N30.00 (F)	Acute cystitis without hematuria
N30.01 (F)	Acute cystitis with hematuria
R10.30 (F)	Lower abdominal pain, unspecified
R30.0 (F)	Dysuria
R30.9 (F)	Painful micturition, unspecified
R31.0 (F)	Gross hematuria
R35.0 (F)	Frequency of micturition
N34.1 (M)	Nonspecific urethritis
N45.1 (M)	Epididymitis
N45.3 (M)	Epididymo-orchitis
N50.811 (M)	Right testicular pain
N50.812 (M)	Left testicular pain
N50.819 (M)	Testicular pain, unspecified

497	Gram Stain (males only) CT and NG symptomatic
A54.01 (M)	Gonococcal cystitis and urethritis
A54.22 (M)	Gonococcal prostatitis
A54.5 (M)	Gonococcal pharyngitis
A54.6 (M)	Gonococcal infection, anus/rectum
A56.01 (M)	Chlamydial cystitis and urethritis
A56.3 (M)	Chlamydial infection, anus/rectum
N34.1 (M)	Nonspecific urethritis
N34.2 (M)	Other urethritis
N45.3 (M)	Epididymo orchitis
N45.1 (M)	Epididymitis
N50.811 (M)	Right testicular pain
N50.812 (M)	Left testicular pain
N50.819 (M)	Testicular pain, unspecified

11361 (CT), 113 or 11363	362 (NG), Chlam	lamydia Trachomatis and Neisseria Gonorrhoeae Additional ICD-10-CM Codes				
	Males and Females		Females Only			
A54.01(M/F)	Gonococcal cystitis and urethritis, unspecified	A54.03 (F)	Gonococcal cervicitis			
A54.5 (M/F)	Gonococcal pharyngitis	N70.03 (F)	Acute salpingitis and oophoritis			
A54.6 (M/F)	Gonococcal infection of anus and rectum	N70.93 (F)	Salpingitis and oophoritis, unspecified			
A56.01 (M/F)	Chlamydial cystitis and urethritis	N72 (F)	Inflammatory disease of cervix uteri			
A56.3 (M/F)	Chlamydial infection of anus and rectum	N89.8 (F)	Vaginal leukorrhea			
A56.4 (M/F)	Chlamydial infection of pharynx	N94.89 (F)	Other conditions assoc with female			
R30.0 (M/F)	Dysuria	N94.10-N94.19 (F)	Dyspareunia			
R30.9 (M/F)	Painful micturition, unspecified		Males Only			
720.2 (M/F)	Contact with or exposure to sexually transmitted	A54.22 (M)	Gonococcal prostatitis	_		
Z20.2 (M/F)	infections	N34.1 (M)	Nonspecific urethritis			
		N34.2 (M)	Other urithritis			
		N45.3 (M)	Epididymo-orchitis			

31532 90887 Females only (age guide)	HPV-amplified probe high-risk (limited to one per year per client) females age 21-99. Additional ICD-10-CM code required beyond ASC-US reflex testing based on age.
D06.9 (≥21)	Carcinoma in situ of cervix
N87.0 (≥21)	Mild cervical dysplasia
N87.1 (≥21)	Moderate cervical dysplasia
R87.610 (≥21)	ASC-US Pap
R87.611 (≥21) R87.612 (≥21)	ASC-H Pap LGSIL Pap
R87.613 (≥21)	HGSIL Pap
R87.616 (≥21)	Satisfactory cervical smear but lacking transformation zone
R87.619 (≥21)	Unspecified abnormal cytological findings, cervix
Z01.42 (≥21)	Pap to confirm normal after abnormal smear
Z87.410 (≥21)	Personal history of cervical dysplasia
•	

509	Hematocrit
Z30.2 (M)	Encounter for sterilization
Z30.09 (F)	Encounter for other general counseling and advice on contraception
Z30.42 (F)	Encounter for surveillance of injectable contraceptive

LAC92214	Wet mount (gel swab)
A59.01 (F)	Trichomonal vulvovaginitis
B37.3 (F)	Candidal vulvovaginitis
N34.1 (M)	Nonspecific urethritis
A59.03 (M/F)	Trichomonal cystitis and urethritis
N34.2 (M)	Other urithritis
N76.0 (F)	Acute vaginitis
Z20.2 (M/F)	Trichomoniasis-exposed partner

3542	Surgical Pathology
A63.0 (M/F)	Anogenital (venereal) warts
B07.9 (M/F)	Viral wart, unspecified
B08.1 (M/F)	Molluscum contagiosum
D06.9 (F)	Carcinoma in situ of cervix
N87.0 (F)	Mild cervical dysplasia
N87.1 (F)	Moderate cervical dysplasia
N88.0 (F)	Leukoplakia of cervix uteri
R87.610 (F)	ASC-US cervical cytology
R87.611 (F)	ASC-H cervical cytology
R87.612 (F)	LGSIL cervical cytology
R87.613 (F)	HGSIL cervical cytology
R87.616 (F)	Satisfactory cervical smear but lacking transformation zone
R87.618 (F)	Other abnormal cytological findings, cervix
R87.619 (F)	Unspecified abnormal cytological findings, cervix
R87.810 (F)	Cervical high-risk HPV positive

3542	Surgical Pathology, LEEP only; females age ≥15 years
D06.9 (F)	Carcinoma in situ of cervix
N87.0 (F)	Mild cervical dysplasia
N87.1 (F)	Moderate cervical dysplasia

Reflex	HPV Genotypes 16, 18/45 females age 30-65 years	
Reflexive from a positive hrHPV test		
R87.810 (F)	Cervical high risk human papillomavirus (HPV) test positive	

19550 or 90521	Thricomonas vaginalis, RNA, TMA Qualitative (female only)
A59.01 (F)	Trichomonal vulvovaginitis
A59.03 (F)	Trichomonal cystitis and urethritis
N76.0 (F)	Acute vaginitis
Z20.2 (F)	Trichomoniasis-exposed partner

For specific indications of diagnostic testing for HPV, CT and NG, and for complication services and corresponding diagnosis codes, please refer to the Family Pact Policies, Procedures, and Billing Instructions Manual, available at medi-cal.ca.gov

Updated March 2024